



# GLOBAL EMERGENCY MEDICAL REGISTRY

## CPR/AED (Adult Ballistic Vest) Emergency First Aid Skills Documentation Form

Candidate (Print): \_\_\_\_\_ Date: \_\_\_\_\_

Examiner (Instructor Name Printed): \_\_\_\_\_

Examiner Signature: \_\_\_\_\_

Evaluation Type:     Initial Certification         Remediation         Renewal

Team Resuscitation:    Team Resuscitation         2 Rescuer         1 Rescuer

Note: Examiner will use a CPR manikin, AED trainer, and filter mask for this skill.

PASS \_\_\_\_\_ FAIL \_\_\_\_\_

Task	Correct	Incorrect
<b><u>Initial Response and Scene Management:</u></b>		
Ensures scene is safe for emergency first aid personnel to approach		
Uses appropriate Personal Protective Equipment (PPE)		
Checks responsiveness appropriately		
Activates emergency medical response or directs bystander to accomplish activation		
<b><u>High Quality Compressions/CPR Section:</u></b>		
Patient wearing body armor and removal is possible, remove the vest immediately in less than 10 seconds		
- If a soft, concealable vest cannot be removed, compressions should be initiated immediately over the vest.		
- If the vest has hard rifle plates, these must be removed or the vest taken off, as they prevent sternal compression	-	
Correct Chest Compression hand placement (center of chest, lower half sternum)		
Chest Compression depth 2–2.4 inches (5–6 cm)		
Chest Compression rate 100–120/min		
Allows full chest recoil		
Minimizes interruptions (<5 seconds) to compressions for ventilation or switch compressors		
Maintains proper body mechanics during chest compressions		
Perform chest compressions continuously until AED, mouth to mask device, and/or second rescuer arrives		
Perform chest compressions correctly at a 30:2 ratio with ventilation once an AED is utilized, a mouth to mask device is available, and/or a second rescuer arrives		
<b><u>Automated External Defibrillator (AED) Operation Section:</u></b>		
Powers on AED immediately		



# GLOBAL EMERGENCY MEDICAL REGISTRY

Applies Defibrillation Pads per manufacturer recommendation		
Pauses CPR and clears all contact with patient during AED rhythm analysis period		
Clears contact with patient before defibrillation shock administered		
Immediately resumes chest compressions after shock is delivered		
<b><u>Airway and Ventilation Section:</u></b>		
Opens airway (head tilt–chin lift or jaw thrust)		
Provides 2 effective ventilations during 30:2 CPR with a one way valve mouth to mask device		
Visible chest rise achieved on patients with normal BMI		
Avoids excessive ventilation		
<b><u>Team Based Resuscitation Section (if applicable)</u></b>		
Communicates clearly with team		
Directs role assignments appropriately		
Coordinates compressor rotation every two minutes with less than 5 seconds of interruption to compressors		
Maintains tactical and safety awareness		

### **Critical Failure Criteria**

Failure to establish oxygenation and compressions (CPR) for the patient
Failure to utilize AED correctly
Failure to identify cardiac arrest
Failure to manage the patient as a competent provider
Exhibits unacceptable affect with patient or other personnel
Uses or orders a dangerous or inappropriate intervention

NOTE: You must factually document any “incorrect” or critical failure criteria on the bottom or back of this form.