

Medication by Auto Injector (EpiPen) Skill Documentation Form

Examiner (Instructor or Licensed Provider):		
Examiner (morried of Licensed Flovider).		
Examiner Signature:		
Pass Fail		
Task:	Correct	Incorrect
Explains procedure to patient, if possible		
Takes Body Substance Isolation procedures and PPE		
Checks for allergies, contraindications, and incompatibilities prior to administration		
Checks Medication for: Correct Medication and dose Solution Clarity Concentration Expiration Date Remove safety cap from autoinjector		
Position injector in the middle of the lateral or anterior thigh		
With one fluid motion, strike the autoinjector into thigh muscle with enough force to activinjection device Hold autoinjector against the thigh long enough to administer medication (10 seconds)	vate	
Quickly withdraw autoinjector		
Dispose of autoinjector into appropriate receptacle		
Obtain a full set of vital signs		
Monitor patient for adverse effects		
Monitor patient for desired effects		
Critical Failure Criteria		
Failure to administer medication within 3 minutes		
Failure to take BSI precautions		
Failure to administer the correct drug, dose, or methodology		
Failure to dispose of needle properly		
Uses inappropriate affect with patient or examiner		
Uses or orders a dangerous or inappropriate intervention NOTE: You must factually document any "incorrect" or critical failure cri		

NOTE: Provide candidate a reclined "Patient" and provide time to familiarize with autoinjector

1

provided for skill.