

Respiratory Case (Reactive Airway Disease) Documentation Form

breathing with rapid respirations; patient's friend states the patient has asthma and uses an inhaler, but the inhaler did not work today and it has been used at least 20 times (If ultrasound BLUE protocol is used, it reveals extensive 'A' line with plural	Candidate (Print):		Date:			
Note: The 70 kg, 16 y/o patient is responsive to voice, able to respond with one word sentences, is in a tripod position and focused on breathing with rapid respirations; patient's friend states the patient has asthma and uses an inhaler, but the inhaler did not work today and it has been used at least 20 times (If ultrasound BLUE protocol is used, it reveals extensive 'A' line with plural movement in all views). The Examiner will use or modify a simulator respiratory case to reflect Sinus Tachycardia at 120 with a MAP of 70 mmHg, SpO2; 8886, ECO2; 55 mmHg with "shark's fin" wavefrom present, RR: 40-50, Temp; 37.5 C. The examiner may conclude the case following proper care by having the patient return to stable with vital signs returning toward stable range. PASS	Exam	niner (Instructor Name Printed):				
breathing with rapid respirations; patient's friend states the patient has asthma and uses an inhaler, but the inhaler did not work today and it has been used at least 20 times (If ultrasound BLUE protocol is used, it reveals extensive 'A' line with plural movement in all views). The Examiner will use or modify a simulator respiratory case to reflect Sinus Tachycardia at 120 with a MAP of 70 mmHg, SpO2: 88%, EtCO2: 55 mmHg with "shark's fin" waveform present, RR: 40-50, Temp: 37.5 C. The examiner may conclude the case following proper care by having the patient return to stable with vital signs returning toward stable range. PASS	Exam	niner Signature:				
Task Correct Incorrect Identifies Respiratory Failure Identifies unstable patient condition Assures proper oxygenation of the patient and prepares for possible intubation Utilizes CPAP or BiPAP to improve ventilation Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Note:	breathing with rapid respirations; patient's friend startoday and it has been used at least 20 times (If ultras movement in all views). The Examiner will use or m MAP of 70 mmHg, SpO2: 88%, EtCO2: 55 mmHg vexaminer may conclude the case following proper care	tes the patient has asthma and uses an inhale sound BLUE protocol is used, it reveals extenodify a simulator respiratory case to reflect with "shark's fin" waveform present, RR: 40	r, but the inl nsive 'A' lin Sinus Tachy -50, Temp:	naler did not wo ne with plural ycardia at 120 v 37.5 C. The	ork vith a
Identifies Respiratory Failure Identifies unstable patient condition Assures proper oxygenation of the patient and prepares for possible intubation Utilizes CPAP or BiPAP to improve ventilation Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel		PASS	Fail			
Identifies unstable patient condition Assures proper oxygenation of the patient and prepares for possible intubation Utilizes CPAP or BiPAP to improve ventilation Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Task			Correct	Incorrect	
Assures proper oxygenation of the patient and prepares for possible intubation Utilizes CPAP or BiPAP to improve ventilation Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Ident	ifies Respiratory Failure				
Utilizes CPAP or BiPAP to improve ventilation Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Identifies unstable patient condition					
Establishes large bore IV above the level of the diaphragm Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Assures proper oxygenation of the patient and prepares for possible intubation					
Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Utilizes CPAP or BiPAP to improve ventilation					
Administers 1-2 gm Magnesium IV Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Estab	olishes large bore IV above the level of the diaphra	agm			
Administers appropriate Steroid IV Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Utilizes Ventolin (Albuterol) nebulization inline with CPAP or BiPAP					
Reassess patient after each intervention and a minimum of every two minutes Note: any "incorrect" represents a skill failure Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Administers 1-2 gm Magnesium IV					
Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Adm	inisters appropriate Steroid IV				
Critical Failure Criteria Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel						
Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Note:	any "incorrect" represents a skill failure				
Failure to establish oxygenation for the patient Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel	Criti	cal Failura Critaria				
Failure to take appropriate medication and/or fluid bolus intervention Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel						
Failure to identify unstable patient Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel			holus intervention			
Failure to manage the patient as a competent provider Exhibits unacceptable affect with patient or other personnel						
Exhibits unacceptable affect with patient or other personnel						
		 				
		•				

NOTE: You must factually document any "incorrect" or critical failure criteria on the bottom or back of this form.

1